

# ERGA Assembly Report

v24.10.15

Tags: ATLASea[ INVALID TAG]

|         |                              |
|---------|------------------------------|
| TxID    | 429176                       |
| ToLID   | <b>xbGlyGlyc1</b>            |
| Species | <i>Glycymeris glycymeris</i> |
| Class   | Bivalvia                     |
| Order   | Arcoida                      |

| Genome Traits     | Expected              | Observed      |
|-------------------|-----------------------|---------------|
| Haploid size (bp) | 3,423,417,414         | 3,667,554,948 |
| Haploid Number    | 14 (source: ancestor) | 19            |
| Ploidy            | 2 (source: ancestor)  | 2             |
| Sample Sex        | Unknown               | Unknown       |

## EBP metrics summary and curation notes

Obtained EBP quality metric for collapsed: 7.8.Q43

The following metrics were automatically flagged as below EBP recommended standards or different from expected:

- Observed Haploid Number is different from Expected
- Kmer completeness value is less than 90 for collapsed
- Assembly length loss > 3% for collapsed

## Curator notes


- Interventions/Gb: 78
- Contamination notes: ""
- Other observations: "The assembly of *Glycymeris glycymeris* (xbGlyGlyc1) is based on 51X ONT data and 204X Arima Hi-C data generated as part of the ATLASea programme (<https://www.atlasea.fr>). The assembly process included the following steps: initial ONT assembly generation with Hifiasm, removal of contaminant sequences using Context, removal of haplotypic duplications using purge\_dups, and Hi-C-based scaffolding with YaHS. In total, 3 contigs were identified as contaminants (bacterial, archaeal, or viral), totaling 0.133 Mb (with the largest being 0.058 Mb). Additionally, 777 regions totaling 272.858 Mb (with the largest being 3.937 Mb) were identified as haplotypic duplications and removed. Mitochondrial genome was assembled using ptGAUL. During manual curation, 66 haplotypic regions were removed, totaling 218Mb (with the largest being 24Mb). Chromosome-scale scaffolds confirmed by Hi-C data were named in order of size. "

## Quality metrics table

| Metrics      | Pre-curation collapsed | Curated collapsed |
|--------------|------------------------|-------------------|
| Total bp     | 3,887,023,287          | 3,667,554,948     |
| GC %         | 34.58                  | 34.59             |
| Gaps/Gbp     | 128.89                 | 136.06            |
| Total gap bp | 50,100                 | 63,300            |
| Scaffolds    | 376                    | 285               |
| Scaffold N50 | 192,936,916            | 189,504,207       |
| Scaffold L50 | 10                     | 9                 |
| Scaffold L90 | 19                     | 18                |
| Contigs      | 877                    | 784               |
| Contig N50   | 14,201,000             | 14,499,000        |
| Contig L50   | 91                     | 84                |
| Contig L90   | 293                    | 265               |
| QV           | 43.0186                | 43.0369           |
| Kmer compl.  | 69.096                 | 66.466            |
| BUSCO sing.  | 89.9%                  | 94.5%             |
| BUSCO dupl.  | 8.3%                   | 3.2%              |
| BUSCO frag.  | 1.2%                   | 1.2%              |
| BUSCO miss.  | 0.7%                   | 1.1%              |


BUSCO: 6.0.0 (euk\_genome\_min, miniprot) / Lineage: mollusca\_odb12 (genomes:36, BUSCOs:4421)

# HiC contact map of curated assembly




collapsed [\[LINK\]](#)

# K-mer spectra of curated assembly




Distribution of k-mer counts per copy numbers found in asm




Distribution of k-mer counts coloured by their presence in reads/assemblies

# Post-curation contamination screening



TAPAs summary Graph

(2 0X contigs have been hidden)



## superkingdom

- Bacteria
- Eukaryota
- N/A

## Longest sequences (bp)

- xbGlyGlyc1\_1 - 237098909 (Eukaryota)
- ▲ xbGlyGlyc1\_2 - 230527397 (Eukaryota)
- xbGlyGlyc1\_3 - 214210711 (Eukaryota)
- + xbGlyGlyc1\_4 - 213863770 (Eukaryota)
- ☒ xbGlyGlyc1\_6 - 197577529 (Eukaryota)

## Length (bp)

- 5.0e+07
- 1.0e+08
- 1.5e+08
- 2.0e+08

**collapsed.** Bubble plot circles are scaled by sequence length, positioned by coverage and GC proportion, and coloured by taxonomy. Histograms show total assembly length distribution on each axis.

## Data profile

| Data     | Long reads | Arima |
|----------|------------|-------|
| Coverage | 51         | 204   |

## Assembly pipeline

```
- Hifiasm
  |_ ver: 0.19.5-r593
  |_ key param: NA
- purge_dups
  |_ ver: 1.2.5
  |_ key param: NA
- YaHS
  |_ ver: 1.2
  |_ key param: NA
```

## Curation pipeline

```
- PretextMap
  |_ ver: 0.1.9
  |_ key param: NA
- PretextView
  |_ ver: 0.2.5
  |_ key param: NA
```

Submitter: Mohamed Meftah Saoues

Affiliation: Genoscope

Date and time: 2026-01-25 11:57:02 CET